

The problem of Initial Data in Cosmology and Conformal General Relativity

Boris Barbashov¹, Piotr Flin^{1,2}, and Victor Pervushin^{1*}

¹ *Bogoliubov Laboratory for Theoretical Physics,*

Joint Institute for Nuclear Research, 141980 Dubna, Russia; and

² *Akademia Swietokrzyska, Instytut Fizyki, ul. Swietokrzyska 15, 25-406 Kielce, Poland*

(Dated: February 8, 2020)

The problem of initial data in cosmology and general relativity is considered by analogy with the Cauchy problem in Newton mechanics where initial data reveal the Galilei symmetry of physical laws, i.e., equations of motion.

We list a set of theoretical and observational arguments (including the latest Supernova data, primordial nucleosynthesis, CMB-temperature, and baryon asymmetry) in favour of conformal symmetry of physical laws in a unified field theory of all interactions, where all dimensional parameters are hidden in masses treated as a scalar (dilaton) field.

PACS numbers: 12.10.-g, 95.30.Sf, 98.80.-k, 98.80.Cq, 98.80.Es, 98.80.Hw

Keywords: General Relativity and Gravitation, Cosmology, Observational Cosmology, Standard Model

Contents

I. Introduction	1
II. Initial Data for the Universe	2
A. Observational cosmology	2
B. Conformal General Relativity	4
C. Relative measurement standard	5
D. Conformal Cosmology	5
E. Conformal version of Dark Matter	7
III. Initial Data in Quantum Cosmology	7
A. Creation of the universe	7
B. Creation of time	8
IV. Fundamental Operator Quantization	9
A. Frame-dependent formulation	9
B. Creation of the matter	10
C. CMB temperature	12
D. Baryon asymmetry	13
V. Summary	14
A. Theory	14
B. Method	14
C. Results: Cold Universe Scenario	15
VI. Conclusion	15
References	16

I. INTRODUCTION

Since the time of Newton, problems in theoretical physics have been reduced to solving differen-

tial equations with the initial data measured by a set of physical devices. "Give me initial data and I explain all world" said Laplace. Initial data were "Great Continental Divide" between theory and experimental physics.

A solution of differential equations in modern theoretical physics also presupposes that some initial data are given. They can be measured by a set of physical instruments which is identified with some frame of reference. Therefore, the dependence of theoretical predictions on the initial data means their frame dependence. In relativistic physics, a frame of reference is determined by the choice of the unit time-like vector, and changes of the frame are fulfilled by its various Lorentz transformations. Until the 60s a system of equations was called the relativistic covariant, if the total space of its solutions for each frame of reference coincided with the total space of its solutions for another one (see [1], Chapter 1, §§1,4,5). Any relativistic system requires the physical meaning only in a definite frame of reference distinguished by the time of axis. On the basis of this definition the frame-dependent fundamental Hamiltonian formulation of relativistic theories was developed [2, 3, 4, 5].

However, by the late 1960s the simplified definition of relativistic covariance as independence of the frame was accepted in theoretical physics as one of the justifications of the frame-free heuristic formulation of relativistic theories such as, for example, the Faddeev-Popov path integral [6]. The idea of the frame-free heuristic formulation belongs to Feynman [7] who noticed that scattering amplitudes of elementary particles in perturbation theory are independent of the reference frame and the gauge choice. The problem of constructing a unitary perturbation theory was posed [8] and solved for non-Abelian theories [6] in the framework of

*e-mail: pervush@thsun1.jinr.ru

the heuristic formulation. In the same approach, the renormalizability of the standard model was proved by Weltman and 't Hooft. The frame-free heuristic formulation actually became only one tool for solving problems of contemporary field theory. However, we should like to note that the applicability of this formulation is restricted to solving scattering problems for elementary particles [9] – the domain where it first appeared [7].

Yet in 1962 Schwinger [4] pointed out that the frame-free formulations can be physically inadequate to the fundamental operator quantization, i.e., they can lose part of information of the initial system or acquire some additional information.

In particular, the frame-dependent formulation of special relativity (SR) by Poincare [10] and Einstein [11] gave the first solution of the problem of energy and time for a relativistic particle in the frame where the time of a particle coincided with its variable X_0 . Barbashov and Chernikov [12] solved the Cauchy problem for a relativistic string in the spirit of the Poincare-Einstein approach [10, 11] choosing a frame where the coordinates on the string were considered as its variables (x_0, x_1) ; in this case, the Born-Infeld theory [12, 13] appears with the resolved problems of time, energy, for any space dimension [14]. The frame-dependent Born-Infeld theory strongly differs from the frame-free formulation with the Virasoro algebra and the problems of time, energy, and space dimension are still unresolved [15]. Thus, we see that Schwinger was right when he rejected all *heuristic* frame-free formulations of relativistic theories "as unsuited to the role of providing the *fundamental* operator quantization" [4].

The frame-dependent *fundamental* formulation of physical theories segregates their frame transformations from the gauge ones.

Frame transformations of variables change a frame and initial data. Historically, the frame symmetries appeared as the Galilei group in the Newton mechanics. The Newton equations are invariant with respect to the Galilei transformations violating initial data of these equations. Generalization of the Galilei group of initial data in relativistic theories is the Poincare group.

Gauge transformations of variables do not alter their initial data and a frame; gauge symmetry establishes constraints of initial data in each frame. The gauge symmetries are principles of construction of modern physical theories and include a group of reparametrizations of coordinates in general relativity.

Modern field theories have their origin in the Faraday-Maxwell electrodynamics. Faraday's profound belief in reality of the electromagnetic fields and the unity of the forces of Nature became the governing principle of the whole further develop-

ment of physics. The strategy of this development was the search for the symmetry principles of electrodynamics and other interactions, whereas the most fruitful tactics for the constructions of practically all working field theories in the 20th century was the analogy with the Faraday-Maxwell electrodynamics; the more we learn about the symmetries of electrodynamics, the more fruitful these analogies are (see in detail [16]). These symmetries are relativistic, gauge, and conformal.

This tactics of the construction of unified theories is the evidence of that the symmetry of the world can be symmetry of electrodynamics, i.e., relativistic, gauge and conformal.

In the present paper, we give a set of arguments showing that physical laws in this world are conformal-invariant, and the conformal symmetry is that of frames of reference themselves. This means that all dimensional parameters violating conformal symmetry are introduced by the initial data in a definite frame of reference.

In Section II, we consider the problem of initial data in observational cosmology and general relativity.

Section III is devoted to the role of the group of reparametrizations of the coordinate time in quantum relativistic cosmology in the resolution of its principal problems of initial data, cosmic singularity, positive arrow of the time measured by our watch, and the status of the Hubble law in quantum theory of the universe.

In Section IV, the frame-dependent formulation of conformal-invariant unified theory is considered to demonstrate its possibilities of explaining origins of matter, Cosmic Microwave Background, and baryon asymmetry of the universe.

II. INITIAL DATA FOR THE UNIVERSE

A. Observational cosmology

The action in Einstein's general relativity (GR)

$$S_{\text{GR}} = - \int d^4x \sqrt{-g} \varphi_0^2 \frac{R(g)}{6} \quad (1)$$

with the Newton constant

$$\frac{\varphi_0^2}{6} = \frac{M_{\text{Planck}}^2}{16\pi} \quad (2)$$

and the measurable interval

$$(ds^2) = g_{\mu\nu} dx^\mu dx^\nu \quad (3)$$

is the basis of the standard cosmological model [17, 18, 19, 20]. This model appears in the homogeneous approximation of the metrics

$$ds^2 = a^2(x^0)[(N_0(x^0)dx^0)^2 - (dx^i)^2], \quad (4)$$

where $a(x^0)$ is the cosmic scale factor, $N_0(x^0)$ is the homogeneous lapse function of the invisible coordinate time x^0 . In observational cosmology this interval is well known as

$$ds^2 = (dt)^2 - a^2(t)(dx^i)^2, \quad (5)$$

here t is the world Friedmann time of the Einstein measurable interval (3)

$$dt = a(\eta)d\eta, \quad (6)$$

and

$$d\eta = N_0(x^0)dx^0 \quad (7)$$

is the conformal time defined as the time of photons flying from any object on the light cone

$$(ds)^2 = (dt)^2 - a^2(t)(dr_c)^2 = 0 \quad (8)$$

with the *coordinate* distance $r_c = \sqrt{x_1^2 + x_2^2 + x_3^2}$. Using the light cone equation $dt = a(t)dr_c$ one can find the coordinate distance - conformal time relation

$$r_c(\eta) = \int_t^{t_0} \frac{dt'}{a(t')} \equiv \eta_0 - \eta, \quad (9)$$

where η_0 is the present-day value of the conformal time when a value of the scale factor is equal to unit $a(\eta_0) = 1$. Therefore, the current cosmological time η of a photon emitted by an atom at the coordinate distance r_c is equal to the difference

$$\eta = \eta_0 - r_c. \quad (10)$$

Observational data testifies that the energy of cosmic photons $E(\eta)$ depends on the coordinate distance (9). The energy of the cosmic photons $E(\eta_0 - r_c)$ (emitted at the conformal time $\eta = \eta_0 - r_c$) is always less then the similar energy of the Earth photons $E_0 = E(\eta_0)$ (emitted at the conformal time η_0):

$$E(\eta) = \frac{E_0}{z(r_c) + 1}, \quad (11)$$

where

$$z(r_c) + 1 = \frac{E_0}{E(\eta_0 - r_c)} \geq 1 \quad (12)$$

is the redshift of the spectral lines E of atoms at objects at the coordinate distance r_c in comparison with the the present-day spectral lines E_0 of atoms at the Earth.

The redshift is explained by varying cosmic scale in the metrics (4) [17, 18, 19, 20]

$$z(r_c) + 1 = a^{-1}(\eta_0 - r_c) \quad (13)$$

with respect to the conformal time with the present-day initial data $a(\eta_0) = 1$.

Fitting observational data (12) with the theory requires the derivation of equations of motion, in particular, the Einstein-Friedmann equation of the balance of the energy densities $\delta S/\delta N_0 = 0$:

$$\frac{\varphi_0^2}{a^2} \left[\frac{da}{dt} \right]^2 = \rho_f(a), \quad (14)$$

in terms of conformal quantities: the conformal time (7), coordinate distance (9), conformal density

$$\rho_c = \rho_f(a)a^4, \quad (15)$$

where $\rho_f(a)$ is the Friedmann density, and other conformal observables ${}^{(n)}F_c$ with the conformal weight (n) , obtained from the Friedmann observables by the conformal transformations

$${}^{(n)}F_c = {}^{(n)}F_f a^{-n}. \quad (16)$$

After these transformations the balance energy equation (14) takes the form [21]

$$\varphi'^2 = \rho_c(\varphi), \quad (17)$$

where $\varphi' = d\varphi/d\eta$, and

$$\varphi(\eta) = \varphi_0 a(\eta) \quad (18)$$

is the running Planck mass.

For the radiation dominant universe $\rho_c = \rho_{\text{rad}} = \text{constant}$ we get a solution of Eq. (17)

$$\varphi(\eta) = \varphi_I \pm \sqrt{\rho_{\text{rad}}}\eta. \quad (19)$$

For the rigid state (when the pressure is equal to energy $p = \rho$)

$$\rho_c = \rho_{\text{rigid}}(\varphi) = \frac{P_\chi^2}{4\varphi^2} \quad (20)$$

a solution of Eq. (17) takes the form

$$\varphi^2(\eta) = \varphi_I^2 \pm P_\chi \eta \equiv \varphi_I^2 [1 \pm 2H_I \eta], \quad (21)$$

where $P_\chi = 2\varphi_I^2 H_I$ is a constant, φ_I, H_I are initial data of φ and its velocity $H = \varphi'/\varphi$ known as the conformal Hubble parameter. This solution can be rewritten in terms of the present-day initial data

$$\varphi^2(\eta) = \varphi_0^2 [1 + 2H_0(\eta - \eta_0)] = \varphi_0^2 [1 - 2H_0 r_c], \quad (22)$$

using the integral of motion $\varphi_I^2 H_I = \varphi_0^2 H_0$, and relation (9) in the case of the increasing scale factor. In this case, the redshift equation (12) takes the form

$$z(r_c) + 1 = \frac{1}{(1 - 2H_0 r_c)^{1/2}}. \quad (23)$$

It gives the coordinate distance – redshift relationship

$$H_0 r_c(z) = \frac{1}{2} \left[1 - \frac{1}{(1+z)^2} \right]. \quad (24)$$

In general case of the sum of a rigid state, radiation, matter (dust) and Λ - term with the partial contributions

$$1 = \Omega_{\text{Rigid}} + \Omega_{\text{Radiation}} + \Omega_{\text{M}} + \Omega_{\Lambda},$$

this coordinate distance – redshift relationship takes the form [22]

$$H_0 r_c(z) = \int_0^z \frac{dz'}{(1+z')^2 \sqrt{\Omega(z')}}}, \quad (25)$$

where

$$\Omega(z') = (1+z')^2 \Omega_{\text{Rigid}} + \Omega_{\text{Radiation}} +$$

$$(1+z')^{-1} \Omega_{\text{M}} + (1+z')^{-4} \Omega_{\Lambda}.$$

Thus, the equation of observational cosmology reveals the possibility to treat GR as a conformal-invariant theory with the world of initial data violating conformal symmetry (like initial data in the Newton equations violate the Galilei symmetry).

B. Conformal General Relativity

Equation (17) follows from the homogeneous approximation of the theory [21, 22, 23, 24], where the Einstein-Hilbert action (1) is changed by the negative Penrose-Chernikov-Tagirov (PCT) action [25]

$$S_{\text{PCT}}[w] = \quad (26)$$

$$\int d^4x \left[\sqrt{-g} w^2 \frac{R(g)}{6} - w \partial_\mu (\sqrt{-g} \partial^\mu w) \right]$$

for the scalar dilaton field w instead of the Planck mass φ_0 with the fixed conformal volume. The equations of motion in this theory do not contain any dimensional parameters; they can be introduced into the theory as nonzero initial data with respect to the conformal time.

The matter fields are described by the Standard Model (SM) with the Higgs potential

$$\mathcal{L}_{\text{Higgs}}[M_{\text{Higgs}}] = -\frac{\lambda}{4} [|\Phi|^2 - M_{\text{Higgs}}^2]^2, \quad (27)$$

where the value of the Higgs mass $M_{\text{Higgs}} = y_{\text{Higgs}} \varphi_0$ can be expressed in the units of the Planck mass (2) $y_{\text{Higgs}} \sim 10^{-17}$. The Higgs mass scales all

masses of elementary particles and it is only one dimensional parameter violating the conformal symmetry of the theory. In fact, the unified theory of all interactions considered as the sum of GR (1) and SM in the Riemannian space:

$$S_{\text{SGR}}^{(1)}[\varphi_0] = S_{\text{SM}}[\varphi_0] + S_{\text{GR}}[\varphi_0] \quad (28)$$

has the hidden conformal symmetry of electrodynamics. This symmetry can be restored if we replace the Planck mass φ_0 by the scalar dilaton field w .

Then, a conformal-invariant version of SGR takes the form [21, 22, 23, 24]

$$S_{\text{CGR}}^{(1)}[w] = S_{\text{SM}}[w] - S_{\text{PCT}}[w], \quad (29)$$

where the Higgs potential (27) becomes

$$\mathcal{L}_{\text{Higgs}}[y_{\text{Higgs}} w] = -\frac{\lambda}{4} [|\Phi|^2 - y_{\text{Higgs}}^2 w^2]^2, \quad (30)$$

and $S_{\text{PCT}}[w]$ is the PCT conformal-invariant action for the dilaton field (26) [25].

The conformal symmetry of the world follows from the mathematical equivalence of SGR (28) to the conformal-invariant theory of the dilaton scalar field, i.e., the conformal general relativity (CGR) (29). This means that all solutions of SGR can be obtained from solutions of CGR by the conformal transformations including the scale ones

$$w^\Omega = \frac{w}{\Omega}; \quad g_{\mu\nu}^\Omega = \Omega^2 g_{\mu\nu}. \quad (31)$$

The dilaton field can be removed from the unified theory (29) by the scale transformation with the scale function

$$\Omega_{\text{gr}} = \frac{w}{\varphi_0} \quad (32)$$

including the dimensional parameter. In this case, the dilaton converts into constant, and CGR (29) coincides with SGR (28)

$$S_{\text{CGR}}^{(1)}[w]|_{w=\varphi_0} = S_{\text{SGR}}^{(1)}[\varphi_0]. \quad (33)$$

This fixing ($w = \varphi_0$) supposes the treatment of the conformal symmetry as a kind of the gauge one. However, the "gauge-fixing" with the dimensional parameter φ_0 introduces an absolute measurement standard violating the scale-invariance of the initial equations of motion. We have seen above that observational cosmology points out another treatment of conformal symmetry as a kind of the frame symmetry. In this case, physical laws in the form of the equations of motion do not contain any dimensional parameters. These parameters can be introduced as initial data violating conformal symmetry like the initial data in the Newton mechanics violate the Galilei symmetry.

Therefore, the way of treating the conformal symmetry as the frame one in the observational cosmology leads to the conformal-invariant and frame-dependent formulation of CGR in terms of conformal quantities *where all dimensional parameters are hidden in masses treated as a scalar dilaton*.

C. Relative measurement standard

It was Weyl's first idea [26] to consider the conformal symmetry as a principle of relativity of all physical measurements. It is clear that we have no absolute physical instruments and have no measurement standards including the Paris meter. If the volume of the universe is expanding, all lengths in the universe are also expanding, including the measurement standard. This means that nothing is expanding, and the measurable spatial volume of the universe is a constant

$$\int d^3x \sqrt{{}^{(3)}g_c} = V_0 = \text{Const.} \quad (34)$$

In this case, the homogeneous approximation keeps the dilaton as a dynamic variable of a running Planck mass

$$w_c(x^0, x^i) = \varphi(x^0). \quad (35)$$

This choice leads to the cosmological equation (17) corresponding to the scale invariance of physical laws at the level of equations of motion in CGR. The scale invariance is broken by initial data like the Galilei invariance of the Newton laws.

Due to the Higgs effect

$$|\Phi| = y_{\text{Higgs}} w(x^0, x^i) = y_{\text{Higgs}} \varphi(x^0) \quad (36)$$

the choice of the relative measurement standard (34), (35) means also running masses of all massive fields of SM.

Therefore, the spectrum of the atom energetic levels (marked by k) is described by the Schrödinger equation

$$\left[\frac{\hat{p}^2}{2m_0 a(\eta)} - \left(\frac{\alpha}{r} + E_k(\eta) \right) \right] \Psi_A = 0. \quad (37)$$

It is easy to check that the exact solution of this equation is expressed through the solution E_k^0 of a similar Schrödinger equation with constant masses m_0 at $a(\eta_0) = 1$

$$E_k(\eta) = a(\eta) E_k^0 \equiv \frac{E_k^0}{z(r_c) + 1}, \quad (38)$$

where $E_k^0 = -m_0 \alpha^2 / k^2$ is the spectrum of a Coulomb atom. It is just above-considered redshift (11) $z(r_c)$ of the spectral lines of atoms at

the coordinate distance $r_c = \eta_0 - \eta$, and η_0 is the present-day value of the conformal time.

The cosmic evolution in the terms of the relative measurement standard (34), (35), and ${}^{(n)}F_c$ can be rewritten in the terms of the absolute standard ${}^{(n)}F_f$ by the conformal transformations (16)

$${}^{(n)}F_c = {}^{(n)}F_f a^{-n} \quad (39)$$

with the scale factor (18)

$$a = \frac{\varphi}{\varphi_0}. \quad (40)$$

Therefore, the absolute measurement standard in the CGR corresponds to the fixation of the measurable volume

$$\int d^3x \sqrt{{}^{(3)}g_f} = a^3 V_0 \quad (41)$$

and the dilaton by constraint

$$w_f(x^0, x^i) = \varphi_0. \quad (42)$$

This choice of variables converts CGR into SGR (33). This means that the homogeneous dilaton $\varphi(\eta)$ in CGR coincides with the scale-factor variable $a(\eta)$ in SGR. Eqs. (34), (35) and (41), (42) are different fixations of measurable variables in CGR marked by F_c and F_f , respectively. These two types of variables are connected by the transformations (39) and (40), including intervals. The conformal interval is the value of the Friedmann interval ds_f in the units of the relative Paris meter a

$$ds_c^2 = a^{-2} ds_f^2. \quad (43)$$

This means that the relative measurement standard identifies the conformal time, coordinate distance and other conformal quantities of the theory with measurable quantities of observational cosmology.

This identification leads to the Conformal Cosmology [22] defined as Standard Friedmann Cosmology in terms of conformal quantities identified with observables, where the Hubble law is treated as the cosmic evolution of all masses [27] including the Planck one, but not the expansion of the universe. A red photon emitted by an atom at a star two billion years (in terms of η) remembers the size of this atom, and after two billion years this photon is compared with a photon of the standard atom at the Earth that became blue due to the evolution of all masses.

D. Conformal Cosmology

We have seen above that the basis of the observational cosmology is the coordinate distance (25) as

a function of the redshift obtained by the equation of a null geodesic conformal interval $(ds_c)^2 = 0$ of light ray traces.

In the comparison with the stationary space (in Standard Cosmology) and stationary masses (in Conformal Cosmology), a part of photons is lost. To restore the full luminosity in both Standard Cosmology (SC) and Conformal Cosmology (CC) we should multiply the coordinate distance by the factor $(1+z)^2$. This factor comes from the evolution of the angular size of the light cone of emitted photons in SC, and from the increase of the angular size of the light cone of absorbed photons in CC [22].

Thus, the redshift - luminosity distance relation in CC is determined by the formula:

$$\ell_{CC}(z) = (1+z)^2 r_c(z). \quad (44)$$

However, in SC we have another definition of measurable distances. They are related to measurable distances in CC (coinciding with the coordinate ones) by the relation

$$r_f(z) = a \int \frac{dt}{a} = \frac{r_c}{1+z}. \quad (45)$$

due to the expansion of the universe. In the units of the absolute Paris meter we have an additional factor $(1+z)^{-1}$. Thus we obtain the relation between two luminosity-distances in SC and CC

$$\ell_{SC}(z) = (1+z)r_c(z) = (1+z)^{-1}\ell_{CC}(z). \quad (46)$$

The absolute Paris meter is the reason of that the latest Supernova data [28, 29] correspond to another equation of state in SC than CC [22].

In Fig. 1 from the paper [22] the predictions of the SC and the CC on the Hubble diagram are confronted with recent experimental data for distant supernovae [28, 29]. Among the CC models the pure rigid state (24) of dark energy with the redshift - luminosity distance relation

$$H_0 \ell_{CC}(z) = z + z^2/2 \quad (47)$$

gives the best description and it is equivalent to the SC fit up to the distance of SN1997ff.

The primordial nucleosynthesis gives us a set of arguments in the favour of the relative measurement standard and conformal-invariance of our world.

The modern data on primordial nucleosynthesis and the element abundance in the universe convincingly testify to that the baryon matter is $30 \div 20$ times smaller than it is necessary for explanation of the evolution of the scale of the universe.

The small contribution of the baryon matter to the evolution of the scale poses the problem on a




FIG. 1: The Hubble diagram [22] for a flat universe model in SC and CC. The points include 42 high-redshift Type Ia supernovae [28] and the reported farthest supernova SN1997ff [29]. The best fit to these data requires a cosmological constant $\Omega_\Lambda = 0.7$ in the case of SC, whereas in CC these data are consistent with the dominance of the rigid state.

source of this evolution in the long epoch of primordial nucleosynthesis (from 10^{-12} up to 10^{11} sec). Its history testifies to the dependence of the scale on observable time as the square root. In the SC this dependence is explained by the dominance of "radiation" with density $\rho_{rad}(a)$. If the baryon matter is insufficient for evolution of the scale, then all known "radiations" are also insufficient for explanations of the evolution of the scale in the epoch of primordial nucleosynthesis. This means that we should find the radiation free explanation of the primordial nucleosynthesis.

Really, the fixed fact of the primordial nucleosynthesis [30] testifies only to that the scale factor is proportional to the square root of the observed time. If we identify the observed time with the conformal one, then the proportionality of the scale factor to the square root of the observed (i.e., conformal) time follows from the rigid state (22).

Therefore, in Conformal Cosmology the square root evolution of the scale factor (22) is compatible with both the latest Supernova data and the primordial nucleosynthesis [30] in agreement with small contribution of radiation in the cosmic evolution (47).

In conformal world the temperature $T_c = T_f a$ is a constant, and we have the mass history [22]

$$m_{era}(z_{era}) = \frac{m_{era}(0)}{(1+z_{era})} \quad (48)$$

with the constant temperature $T = 2.73$ K = 2.35×10^{-13} GeV where $m_{era}(0)$ is characteristic energy (mass) of the era of the universe evolution, which begins at the redshift z_{era} .

Eq. (48) has an important consequence that all physical processes, which concern the chemical composition of the universe and depend basically on the Boltzmann factors with the argument

$$\frac{m_c(z)}{T_c(0)} = \frac{m_f(0)}{T_f(z)}, \quad (49)$$

cannot distinguish between the Conformal Cosmology (m_c/T_c) in the rigid state with the square root dynamics and the Standard Cosmology (m_f/T_f) in the radiation state with the same square root dynamics in the "absolute" measurement standard. From this formula it is clear that the z -history of masses with invariant temperatures in the rigid state of Conformal Cosmology is equivalent to the z -history of temperatures with invariant masses in the radiation stage of the Standard Cosmology.

Thus, the description of eras in Conformal Cosmology [22] with the relative measurement standard strongly differs from the one in Standard Cosmology.

Standard Cosmology describes the Hot Universe Scenario [31] with three eras: inflationary, radiation, and again inflationary *quintessence*, and with unsolved questions at the level of SGR:

1. Why is the radiation-dominance era changed again by the inflationary *quintessence*, as it follows from the latest Supernova data?

2. How can the appearance of the pure radiation era between two inflationary ones, the fast – primordial and the slow – *quintessence*, be explained?

and two more questions at the level of CGR:

3. What is a physical origin of the Planck era of the Early Universe with initial data $a_I \simeq 10^{-60}$, if the Planck scale belongs to the present-day value of the dilaton?

4. What is a physical origin of the absolute measurement standard (the Paris meter) in the expanding universe?

After the conformal transformation (16) of Standard Cosmology we obtain Conformal Cosmology with the Cold Universe Scenario where the last two eras (the chemical evolution, and the present-day cosmic evolution of all masses) are described by the single equation of the rigid state. In the conformal world the conformal volume and temperature are not varying. The CMB temperature $2.7K$ becomes the fundamental parameter. The SC initial data as a Hot Fireball of massless particles (undergoing a set of phase transitions) are forbidden in quantum field theory due to the mass singularity of massive vector bosons [32, 33]. Therefore, new questions appear:

1. What is an origin of the rigid state?
2. What is an origin of the CMB temperature?
3. What is a primordial state of matter admitted by CGR at the level of quantum field theory?

4. What are reasons for the creation of the universe, its time, and matter with the CMB radiation and the baryon asymmetry?

E. Conformal version of Dark Matter

The dominance of the rigid state in the cosmic evolution of the scale factor means that this cosmic evolution is described by the action for a *quintessence* scalar field χ

$$S_{\text{CGR}}^{(2)}[w] = S_{\text{CGR}}^{(1)}[w] + S_Q[w, \chi] \quad (50)$$

where

$$S_Q[\chi] = \int d^4x \sqrt{-g} w^2 \partial_\mu \chi \partial^\mu \chi. \quad (51)$$

This action appears in the bi-dilaton theory [34] defined as a difference of two dilaton actions (26)

$$S_{\text{PCT}}[w_-] - S_{\text{PCT}}[w_+] = \quad (52)$$

$$-S_{\text{PCT}}[w] + S_Q[w, \chi]$$

for $w_+ = w \cosh \chi$ and $w_- = w \sinh \chi$. At the level of SGR (obtained from CGR by the substitution $w = \varphi_0$) this means that we add a massless scalar field

$$S_{\text{SGR}}^{(2)}[\varphi_0] = S_{\text{CGR}}^{(1)}[\varphi_0] + S_Q[\varphi_0, \chi] \quad (53)$$

which is not interacting with other fields except for gravitational ones.

In the homogeneous approximation the scalar part of this action takes the form

$$S = V_0 \int_{x_1^0}^{x_2^0} \frac{dx^0}{N_0} [-(\partial_0 \varphi)^2 + \varphi^2 (\partial_0 \tilde{\chi})^2]. \quad (54)$$

This system has an integral of motion of the rigid state ($\rho = p$)

$$P_\chi = 2\varphi^2 \tilde{\chi}' = \pm (\varphi^2)'. \quad (55)$$

The rigid state of *quintessence* χ leads to the square root evolution of the scale factor, or the running Planck mass (22)

$$\varphi^2(\eta) = \varphi_I^2 + P_\chi \eta \equiv \varphi_I^2 (1 \pm 2|H_I| \eta). \quad (56)$$

III. INITIAL DATA IN QUANTUM COSMOLOGY

A. Creation of the universe

The dynamic of the cosmic scale factor with respect to the conformal time is determined in GR in

homogeneous approximation (35), and (43). This approximation gives the relativistic cosmology

$$S = -V_0 \int_{x_1^0}^{x_2^0} dx^0 \left[\frac{1}{N_0} \left(\frac{d\varphi}{dx^0} \right)^2 + N_0 \rho(\varphi) \right] \quad (57)$$

in the conformal flat space (43) inheriting the group of symmetry of GR in the form of invariance with respect to reparametrization of the coordinate time $x^0 \rightarrow \tilde{x}^0 = \tilde{x}^0(x^0)$.

These models with different energy densities $\rho(\varphi)$ (described firstly at the end of the 60s by DeWitt, Wheeler, and Misner [35]) directly correspond to SR [14, 36, 37, 38]

$$S_{\text{SR}} = -\frac{m}{2} \int_{\tau_1}^{\tau_2} d\tau \left[-\frac{\dot{X}^{\mu 2}}{e} + e \right]. \quad (58)$$

In action (57), the conformal time $N_0 dx^0 = d\eta$ plays the role of the geometric interval $ed\tau = ds$; the cosmic scale factor (or dilaton) φ , the dynamic evolution parameter X_0 ; and its canonical momentum P_φ , the dynamic energy of the universe $P_0 = \sqrt{P_\varphi^2 + m^2}$.

The Hamiltonian form of the action (57)

$$S = \int_{x_1^0}^{x_2^0} dx^0 \left[-P_\varphi \partial_0 \varphi + \frac{N_0}{4V_0} (P_\varphi^2 - E^2) \right], \quad (59)$$

where $E = 2V_0 \sqrt{\rho(\varphi)}$ gives the well known Wheeler-DeWitt quantum equation in the field space

$$(P_\varphi^2 - E^2(\varphi)) \Psi_{\text{field}}[\varphi] = 0. \quad (60)$$

In particular, in the case of radiation $\rho_c = \text{const.}$, the universe wavefunction $\Psi_{\text{field}}[\varphi]$ is the sum over two values of the dynamic energy $P_\varphi = \pm E$,

$$\Psi_{\text{field}}[\varphi] = A_E^+ e^{is(\varphi_0, \varphi_I)} + A_E^- e^{-is(\varphi_0, \varphi_I)}, \quad (61)$$

where $s(\varphi_0, \varphi_I) = [\varphi_0 - \varphi_I]E$, and A_E^+ is interpreted as the operator of creation of a universe and A_E^- is interpreted as the operator of annihilation of an antiuniverse. Such an interpretation (similar to the Dirac causality quantization of the relativistic particle) removes negative energies and associates positive frequencies with the forward motion ($P_\varphi > 0$, $\varphi_0 > \varphi_I$) and negative frequencies with the backward motion ($P_\varphi < 0$, $\varphi_0 < \varphi_I$). Here φ_0 is the present-day value of the dilaton (i.e., scale factor) and φ_I is the initial data of the dilaton as the moment of the creation of the universe in the world space.

Thus, we can see that the relativistic cosmology (57) can explain not only the Hubble law, but also a creation of the universe without its cosmic singularity, as for a positive energy $E \geq 0$ the current φ is always greater than φ_I .

The similar wavefunction for the rigid state was constructed in [38].

B. Creation of time

The symmetry of the theory with respect to reparametrizations of the coordinate time x^0 means that this time is invisible.

As it was shown in [14, 36, 38], the passage to the visible geometric interval $d\eta = N_0 dx^0$ as dynamic variable can be accomplished with a Levi-Civita (LC) - type canonical transformation

$$\varphi, P_\varphi \implies Q_0, \Pi_0 \quad (62)$$

turning energy into a new momentum and the scale factor into new variable Q_0 coinciding with the conformal time η . In particular, for the radiation universe (61) the L-C transformation takes a form

$$P_\varphi = \pm 2\sqrt{V_0 \Pi_0}; \quad \varphi = \pm \sqrt{\frac{\Pi_0}{V_0}} Q_0. \quad (63)$$

After this transformation the action (59) becomes

$$\int_{x_1^0}^{x_2^0} dx^0 [-\Pi_0 \partial_0 Q_0 + N_0 (\Pi_0 - V_0 \rho) + \partial_0 (\Pi_0 Q_0)].$$

The variation of the action with respect to the momentum Π_0 gives the equation $dQ_0 = N_0 dx^0 \equiv d\eta$. The variation with respect to the lapse function N_0 gives the energy constraint $\Pi_0 - V_0 \rho = 0$. The quantum equation for wavefunction takes the form

$$\frac{d}{id\eta} \Psi_{\text{geometric}}[\eta] = V_0 \rho_c \Psi_{\text{geometric}}[\eta]. \quad (64)$$

The explicit form of this wavefunction in the rigid state is given in [38]. Thus, a relativistic universe (like a relativistic particle in SR) has two physical *realities*: the *field* and *geometric*. Each *reality* has its measurable evolution parameter (φ , or η), nonzero Hamiltonian, energy, and wavefunction ($\Psi_{\text{field}}[\varphi]$, or $\Psi_{\text{geometric}}[\eta]$).

The Hubble law $\varphi(\eta)$, i.e. the cosmic evolution of the scale factor φ on the geometric time η , in the form of the coordinate-distance – redshift relation is the relation of two evolution parameters of two physical *realities*. The Hubble law is the pure relativistic effect of the type of the relativistic shrinking in SR.

The quantum stability of the system in the *field* space reached by the Dirac causality quantization for the universe results in the positive arrow of geometric time $\eta \geq 0$. Therefore, the quantum relativistic cosmology describes the Hubble with the initial data $\varphi(\eta = 0) = \varphi_I$ of creation of the universe with the positive energy $E = P_\varphi$ at the point φ_I in the *field* space with the positive arrow of time η in the *geometric* space.

IV. FUNDAMENTAL OPERATOR QUANTIZATION

A. Frame-dependent formulation

The next problem of quantum creation of matter exceeds the limits of the homogeneous approximation. To consider this problem, we need the formulation of CGR at the level of quantum field theory, i.e., Quantum Gravity. Let us consider Quantum Gravity on the basis of the action $S_{\text{CGR}}^{(2)}[w|F]$ given by (50) in the world space of all field variables $[w|F]$. This action reflects the group of diffeomorphisms of a complete set of all initial data in the conformal world, measured by a set of physical instruments in the units of measurement standards. It is worth reminding that J.C. Maxwell wrote: "The most important aspect of any phenomenon from mathematical point of view is that of a measurable quantity. I shall therefore consider electrical phenomena chiefly with a view to their measurement, describing the methods of measurement, and defining the standards on which they depend" [39].

Measurable quantities and standards are determined in a definite frame of reference. The frame-dependent *fundamental operator quantization* begins with the definition of a frame of reference of the Hamiltonian description [40, 41, 42]

$$ds^2 = (Ndx^0)^2 - \quad (65)$$

$${}^{(3)}g_{ij}(dx^i + N^i dx^0)(dx^j + N^j dx^0)$$

and its group of diffeomorphisms [43] including reparametrizations

$$x^0 \rightarrow \tilde{x}^0 = \tilde{x}^0(x^0) \quad (66)$$

and gauge transformations

$$x^i \rightarrow \tilde{x}^i = \tilde{x}^i(x^0, x^i). \quad (67)$$

We have seen in the previous sections that the reparametrization symmetry (66) means that the coordinate parameter x^0 is invisible, and its Hamiltonian of evolution is equal to zero. The visible

time of the *fundamental operator quantization* is obtained by the Levi-Civita canonical transformation [36] of the variable with a negative contribution to this Hamiltonian. It is just the dilaton.

Therefore, the identification of the dilaton with a dynamic evolution parameter by (34) and (35) is not the homogeneous approximation, but the fixation of a frame in the field world space ($w|F$) of CGR is in agreement with the method of solving the problem of energy and time in relativistic theories proposed by Poincare and Einstein (PE) in SR [10, 11], and by Barbashov and Chernikov (BC) in the theory of a string [12, 13, 14]. After the PE-BC fixation of a frame the action (50) takes the form

$$S_{\text{CGR}}^{(2)}[w|F] = S_{\text{SGR}}^{(2)}[\varphi|F_c] + S_{\text{frame}}[\varphi], \quad (68)$$

where $S_{\text{SGR}}^{(2)}[\varphi|F_c]$ is the SGR with the running Planck mass and

$$S_{\text{frame}}[\varphi] = \int_{x_1^0}^{x_2^0} dx^0 \int_{V_0} d^3x \varphi \frac{d}{dx^0} \left(\sqrt{{}^{(3)}g_c} \frac{d\varphi}{N_c dx^0} \right) + \int_{x_1^0}^{x_2^0} dx^0 \Lambda(x^0) [V[g_c] - V_0], \quad (69)$$

follows from the PCT action of the dilaton, and the last term with the Lagrange multiplier $\Lambda(x^0)$ gives the constraint of the constant relative volume

$$V[g_c] = \int d^3x \sqrt{{}^{(3)}g_c}. \quad (70)$$

Remind that in quantum theory of gravity instead of (68), (69) one conventionally uses the frame-free expression (33), (53) [40, 41, 42]

$$S_{\text{CGR}}^{(2)}[w|F]|_{(w=\varphi_0, F=F_f)} = S_{\text{SGR}}^{(2)}[\varphi_0|F_f], \quad (71)$$

where fields F_f are treated as measurable quantities. The theory (71) expressed in terms of the conformal quantities F_c , (39) and $\varphi = \varphi_0 a$ (40) becomes equivalent to (68), (69) [38].

However, if we forget about $S_{\text{frame}}[\varphi]$ in (68), considering the absolute variables F_f as measurable quantities in (71) [24, 42], we get the frame-free theory that differs from the frame-dependent one (68) like the frame-free Virasoro algebra dynamics [15] differs from the frame-dependent Born-Infeld theory of a string [12, 13, 14].

The frame-free formulation (71) loses, together with $S_{\text{frame}}[\varphi]$, the internal evolution parameter φ and the invariant geometric time $d\eta = N_0(x^0)dx^0$, where N_0 is the global lapse function defined in

$S_{\text{frame}}[\varphi]$ by the Gibbs averaging the relative lapse function N_c ,

$$N_0^{-1} = \langle N_c^{-1} \rangle \equiv \frac{\int_{V_0} N_c^{-1} \sqrt{{}^{(3)}g_c} d^3x}{\int_{V_0} \sqrt{{}^{(3)}g_c} d^3x}. \quad (72)$$

This means that the frame-free formulation (71) loses the dynamic energy of the universe as a canonical momentum of the dilaton in the Hamiltonian approach to the theory (68). In this approach, the action (68) takes the form

$$S_{\text{CGR}}[\varphi|F_c] = \int dx^0 \left\{ \left[\int d^3x \sum_{F_c} P_F \partial_0 F_c \right] - \right. \quad (73)$$

$$\left. P_\varphi \partial_0 \varphi + N_0 \left[\frac{P_\varphi^2}{4V_0} - V_0 \rho_{\text{tot}} \right] + \Lambda(V - V_0) \right\};$$

here the relative Hamiltonian $V_0 \rho_{\text{tot}} = H_{\text{tot}}[\varphi|F_c]$ is the conventional one [40, 41, 42]

$$V_0 \rho_{\text{tot}} = \int d^3x \left[\mathcal{N}\mathcal{H} - N^k \mathcal{P}_k + C_0 P_g + C_i f^i \right],$$

where all fields are changed by the relative one, and the Planck mass φ_0 is changed by the running Planck mass, $N_0[g_c]$, and $V[g_c]$ are considered as functionals given by eqs. (72) and (70), $\mathcal{N} = N_c/N_0$, N^k , C , C^k are the Lagrangian multipliers for the constraints including the Dirac conditions of transverseness $f^j = \partial_i [g_c^{ij} |g_c|^{1/3}] = 0$ and the minimum embedding of three-dimensional hypersurface into the four-dimensional Riemannian space-time with the zero momentum of spatial metric determinant $P_{g_c} = 0$ [40].

The frame-dependent GR is the direct field generalization of the frame-dependent SR (58). We have the one - to - one correspondence between SR and GR [14, 36, 37], i.e., their proper times (i.e., geometric intervals)

$$ds = ed\tau \quad \iff \quad d\eta = N_0 dx^0, \quad (74)$$

their world spaces

$$X_0, X_i \quad \iff \quad \varphi, F_c, \quad (75)$$

their energies

$$P_0 = \pm \sqrt{P_i^2 + m^2} \quad \iff \quad P_\varphi = \pm 2V_0 \sqrt{\rho_{\text{tot}}}, \quad (76)$$

and their two-time relations in the differential form

$$\frac{dX_0}{ds} = \pm \frac{\sqrt{P_i^2 + m^2}}{m} \quad \iff \quad \frac{d\varphi}{d\eta} = \pm \sqrt{\rho_{\text{tot}}} \quad (77)$$

and in the integral forms

$$s = \pm \frac{m}{\sqrt{P_i^2 + m^2}} X_0 \quad \iff \quad \eta = \pm \int_{\varphi_I}^{\varphi_0} \frac{d\varphi}{\sqrt{\rho_{\text{tot}}}}. \quad (78)$$

The equations of motion of this theory were derived in [14, 38].

The observational data testify to that the total energy density

$$\rho_{\text{total}} = \rho_{\text{rigid}}(\varphi) + H_{\text{qft}}/V_0 \quad (79)$$

contains a large homogeneous part $\rho_{\text{rigid}}(\varphi)$ and small local excitations described in the lowest order of perturbations by the Hamiltonian of the evolution of the free fields $H_{\text{qft}} = H_{\text{qft}}^{(2)} + H_{\text{int}}$. In this case, the first order of the series of the action

$$P_\varphi d\varphi = 2V_0 \varphi_0 \sqrt{\rho_{\text{rigid}}(\varphi)} da + H_{\text{qft}} d\eta + O(\delta^2);$$

in $H_{\text{qft}}/V_0 \rho(\varphi) = \delta$ coincides with the action of quantum field theory in terms of the conformal quantities ${}^{(n)}F_c = {}^{(n)}F a^{-n}$; here equation (77) is used to get $d\eta \simeq d\varphi/\sqrt{\rho_{\text{rigid}}(\varphi)}$. Therefore, the Quantum Gravity in the approximation under consideration is the sum of Quantum Cosmology considered above and quantum field theory in terms of the conformal quantities.

We have seen above that two realities of the universe in the conformal world solved the problems of creation of the quantum universe without the cosmic singularity and creation of its time with a positive arrow.

We left for consideration the mystery of the quantum creation of matter from vacuum.

B. Creation of the matter

The problem of cosmological particle creation in strong gravitational fields, in particular in the vicinity of the cosmological singularity, has been topical for more than thirty years [44, 45, 46, 47, 48, 49]. It is considered that the number of created particles is obviously not sufficient for the description of primordial element abundance in the early universe [46]. Therefore, in the inflationary models [31] it is proposed that from the very beginning the universe is a hot fireball of massless particles that undergo a set of phase transitions. However, the origin of particles is an open question as the isotropic evolution of the universe cannot create massless particles [44, 45, 46, 47, 48, 49]. Nowadays, it is evident that the problem of the cosmological creation of matter from vacuum is beyond the scope of the inflationary model.

The investigation of the cosmological creation of massive vector bosons from vacuum [34, 50, 51]

introduced corrective amendments to the modern model of the isotropic evolution of the universe, since vector bosons detected at the CERN accelerator are unique particles of the Standard Model (SM) which can exhibit a space singularity [32, 33].

The matter is that the phenomenon of the cosmological creation in both Standard Cosmology and Conformal one is described by the diagonalization of equations of motion in terms of conformal fields and coordinates. In terms of conformal quantities (as we have seen above) the cosmic scale factor is the scale of all masses, including the Planck mass and masses of particles in the field theories, in particular in SM. For massive conformal fields the cosmological singularity means disappearance of masses. The massive vector bosons are unique particles of SM which have a mass singularity [33]. The absence of the massless limit of the massive Yang-Mills theory is a well-known fact [32]. It leads to an ultra-violet divergence of the number of created longitudinal bosons calculated in the lowest order of perturbation theory [50, 51]. In the papers [34, 50, 51] theoretical and observational arguments were listed in favour of the consideration of created vector bosons as primary particles whose decay products form all visible matter of the universe, including CMBR and galaxies with baryon asymmetry.

Let us consider "free" vector W^\pm, Z^0 bosons $v_\mu = (\mathbf{v}_0, \mathbf{v}_i)$ in the conformal-flat metric

$$ds_c^2 = (\bar{N}_0(x^0)dx^0)^2 - (dx^i)^2. \quad (80)$$

with the cosmic evolution of their mass $M_v a = y_v \varphi$ in the rigid state, where $a = \varphi/\varphi_0$. Massive vector bosons in SM are described by the Lagrangian density

$$\mathcal{L}_v = -\frac{F_{\mu\nu}F^{\mu\nu}}{4} - \frac{M_v^2 a^2 v_\mu v^\mu}{2},$$

where $F_{\mu\nu} = (\partial_\mu v_\nu - \partial_\nu v_\mu)$. Their mass singularity is revealed by the fundamental operator quantization [3, 4, 33].

This quantization is formulated in a definite frame of references where the time component v_0 of a vector field is distinguished. This component has zero canonical momentum $p_0 = \partial\mathcal{L}/\partial(\partial_0 v_0) = 0$. The condition of conservation of the zero momentum is the classical equation $\delta S/\delta v_0 = 0$

$$\left[\bar{\partial}^2 - (M_v a)^2\right] v_0 = \partial_0 \bar{\partial} \vec{v} \quad (81)$$

As both the field and its momentum are fixed by the constraints the quantization of this field contradict to the uncertainty principle. Therefore, the time component is treated as pure classical one, and it is excluded from the action by resolving the classical equation (81) before quantization. The relativistic covariance of the procedures

of excluding of the time component and operator quantization of the spatial ones was proved in [33]. The results of this paper can be presented in the Hamiltonian form of the total action (including the dilaton part) in terms of the Fourier-components

$$\mathbf{v}_k = \int_{V_0} d^3x e^{i\mathbf{k}\mathbf{x}} \mathbf{v}(\mathbf{x})$$

$$S_{\text{tot}} = \int_{x_1^0}^{x_2^0} dx^0 \left(\sum_k \left[\mathbf{p}_k^\perp \partial_0 \mathbf{v}_k^\perp + \mathbf{p}_k^\parallel \partial_0 \mathbf{v}_k^\parallel \right] - P_\varphi \partial_0 \varphi \right) + \int_{x_1^0}^{x_2^0} dx^0 N_0 \left[\frac{P_\varphi^2}{4V_0} - V_0 \rho_{\text{tot}} \right],$$

where

$$\rho_{\text{tot}} = \frac{P_\chi^2}{4\varphi^2} + \rho_v, \quad (82)$$

$$\rho_v = V_0^{-1}(H^\perp + H^\parallel); \quad (83)$$

H^\perp and H^\parallel represent Hamiltonians of free fields for transverse and longitudinal components of vector bosons

$$H^\perp = \sum_{k,\sigma} \frac{1}{2} \left[\mathbf{p}_k^{\perp 2} + \omega^2(\varphi, k) \mathbf{v}_k^{\perp 2} \right], \quad (84)$$

$$H^\parallel = \sum_{k,\sigma} \frac{1}{2} \left[\left(\frac{\omega(\varphi, k)}{M_v a} \right)^2 \mathbf{p}_k^{\parallel 2} + (M_v a)^2 \mathbf{v}_k^{\parallel 2} \right]$$

with the dispersion relation in the form of the one particle energy $\omega(\varphi, k) = \sqrt{\mathbf{k}^2 + (y_v \varphi)^2}$, $a = \varphi/\varphi_0$, and $P_\chi = 2\varphi_0^2 H_0 = 2\varphi_I^2 H_I$ is integral of motion of the rigid state (20).

In field theories "particles" are defined as field variables a^\pm , a for which their Hamiltonian takes the form of production of the one-particle energy and the operator of number of particles $\hat{N}^J = a_\sigma^{J+}(-k, x^0) a_\sigma^J(k, x^0)$:

$$H^J = \sum_{k,\sigma,J} \omega(\varphi, k) \hat{N}_{k,\sigma}^J,$$

where σ is the spin, and $J = \perp, \parallel$ designate transverse and longitudinal bosons. In particular, the definition of "particles" for longitudinal bosons takes the form

$$\mathbf{v}_k^\parallel = \frac{\sqrt{\omega(\varphi, k)}}{y_v \varphi \sqrt{2V_0}} (a^+(-k) + a^\parallel(k))$$

$$\mathbf{p}_k^\parallel = -i \frac{y_v \varphi}{\sqrt{2V_0 \omega(\varphi, k)}} (a^{\parallel+}(-k) - a^\parallel(k)).$$

We can see that this definition of quantum particles contains the mass singularity $M_v a =$

$y_v \varphi(\eta) \rightarrow 0$. The dependence of masses on time leads to nondiagonal terms in the canonical differential form [37, 50, 51]:

$$p_k \partial_0 v_k = \frac{i}{2} [a^+(k) \partial_0 a(-k) - a(k) \partial_0 a^+] - \frac{i}{2} [a^+(k) a^+(-k) - a(k) a(-k)] \partial_0 \Delta_k$$

where

$$\Delta_v^\perp(\varphi) = \frac{1}{2} \ln \left(\frac{\omega_v}{\omega_I} \right),$$

$$\Delta_v^\parallel(\varphi) = \ln \left(\frac{\varphi}{\varphi_I} \right) - \frac{1}{2} \ln \left(\frac{\omega_v}{\omega_I} \right),$$

and φ_I and ω_I are cosmic initial data as a point of the creation of the universe in the world field space. Just these nondiagonal terms are sources of the cosmological creation of particles. The classical equations in terms of "particles" become diagonal $(b^+)' = -i\omega_b b^+$, $b' = i\omega_b b$ after the Bogoliubov transformations to "quasiparticles"

$$b_\zeta = \alpha_\zeta a_\zeta + \beta_\zeta a_\zeta^+ \quad (85)$$

with the Bogoliubov energy ω_b . These transformations playing the role of the Levi-Civita ones from the *field* set of variables (φ, a^+, a) to the *geometric* set (η, b^+, b) [36].

The coefficients $|\beta|$ determine the number of particles

$$\mathcal{N}_\zeta(\eta) = {}_{\text{sq}}\langle 0 | \hat{N}_\zeta | 0 \rangle_{\text{sq}} = |\beta_\zeta|^2 \quad (86)$$

created during the time η from "squeezed" vacuum defined as

$$b_\zeta |0\rangle_{\text{sq}} = 0. \quad (87)$$

The density of created particles is

$$\rho_v = \sum_\zeta \omega_\zeta(\varphi) {}_{\text{sq}}\langle 0 | \hat{N}_\zeta | 0 \rangle_{\text{sq}}. \quad (88)$$

The Bogoliubov equations of the diagonalization rewritten in the dimensionless variables $\tau = \eta 2H_I = \eta/\eta_I$, $x = q/M_I$ and initial data M_I , H_I with taking into account the dispersion relation $\omega_v = M_I \sqrt{1 + \tau + x^2}$, were solved numerically at positive values of momentum $x = q/M_I$, by utilizing the asymptotics of the solutions $r(\tau) \rightarrow \text{const} \cdot \tau$, $\theta(\tau) = \pi/4 + O(\tau)$ from the neighborhood $\tau = 0$. The distribution functions of the longitudinal $\mathcal{N}^\parallel(x, \tau)$ and transverse $\mathcal{N}^\perp(x, \tau)$ vector bosons for the initial data $H_I = M_I$, are introduced in Fig. 2.

The choice of these initial data is determined by the lower boundary for a boson mass from the area of its initial values allowed by the uncertainty principle $\delta E \eta_I \geq 1$ for variations of energy $\delta E = 2M_I$ at creation of a pair of bosons in the universe with minimum lifetime for a considered case $\eta_I = 1/2H_I$. Consequently, we get $M_I \geq H_I$.




FIG. 2: Dependence of longitudinal N^\parallel and transverse N^\perp components of the distribution function of vector bosons from dimensionless time $\tau = 2H_I\eta$ and dimensionless momentum $x = q/M_I$ at value of the initial data $M_I = H_I$ [50, 51].

C. CMB temperature

From Fig. 2. one can see that a longitudinal component of the distribution function is essentially larger than the transverse one, which demonstrates a more intensive cosmological creation of

longitudinal bosons in contrast with transverse ones. The sluggish decrease with momentum of longitudinal components is explained by mass singularity of a distribution function of longitudinal vector bosons [32, 33]. One of the consequences of such a decrease is the divergence of the density of created particles [45]

$$n_v(\eta) = \frac{1}{2\pi^2} \int_0^\infty dq q^2 \left[\mathcal{N}^\parallel(q, \eta) + 2\mathcal{N}^\perp(q, \eta) \right] \rightarrow \infty.$$

This divergence is the shortage of the lowest order of perturbation theory, where one neglects interactions of vector bosons, including the scattering processes forming an integral of collisions in the kinetic equation for distribution functions.

The thermal equilibrium is formed, if the relaxation time of the process of establishment of temperature of vector bosons

$$\eta_{\text{relax.}} = [n(T^3)\sigma_{\text{scat.}}]^{-1} \quad (89)$$

is less than the time for creating vector bosons. Fig. 2. shows us that the time of establishment of the number of bosons is the order of lifetime of the Universe $\eta_I = 1/2H_I$. This means that

$$n(T^3)\sigma_{\text{scat.}} \sim H_I. \quad (90)$$

The dominating contribution of large momenta to the integral of the density of the particle number (see Fig. 2.) means the relativistic dependence of the density on temperature

$$n_v \sim T^3.$$

Using that $\sigma_{\text{scat.}} \sim 1/M_I^2$ one can estimate this temperature from (90) as an integral of motion of the rigid state

$$T \sim [M_I^2 H_I]^{1/3} = [M_0^2 H_0]^{1/3} \simeq 2.7K. \quad (91)$$

Remind that the rigid state in the conformal cosmology is compatible with the latest Supernova data and the z-history of the chemical evolution of the element abundance [22], therefore we can utilize the rigid equation of state for estimation of temperature of the CMB radiation as a final product of the decays of primordial vector bosons.

On the other hand, it is possible to estimate the lifetime of the created bosons in the early universe in dimensionless unities $\tau_L = \eta_L/\eta_I$, where $\eta_I = (2H_I)^{-1}$, by utilizing an equation of state $a^2(\eta) = a_I^2(1 + \tau_L)$ and define the lifetime of W -bosons in the Standard Model for $M_I = H_I$

$$1 + \tau_L = \frac{2 \sin^2 \theta_W}{\alpha a(\eta_L)} = \frac{2 \sin^2 \theta_W}{\alpha \gamma_v \sqrt{1 + \tau_L}}, \quad (92)$$

where θ_W is the Weinberg angle, $\alpha = 1/137$.

The solution of equation (92)

$$1 + \tau_L = \left(\frac{2 \sin^2 \theta_W}{\alpha} \right)^{2/3} \simeq 16 \quad (93)$$

gives an estimation of the lifetime of the created bosons $\tau_L \simeq 15$ which is 15 times larger than the relaxation time. Therefore, we can introduce the concept of the temperature of vector bosons, which is inherited by final products of their decays, i.e. gamma-quanta, forming, according to the modern point of view, the CMB radiation in the universe.

The temperature of photon radiation, which appears after annihilation and decays of W^\pm and Z bosons in the conformal cosmology, is invariant and the simple estimation fulfilled above gives the value surprisingly close to the observed temperature of the CMB radiation which is determined in the conformal cosmology as a fundamental constant - the integral of the rigid state (91).

The back reaction of created particles on the evolution of the Universe is very small. The equation of motion $\varphi'^2(\eta) = \rho_{\text{tot}}(\eta)$ with the Hubble parameter defined as $H = \varphi'/\varphi$ means that the energy density of the universe at any moment is equal to the so-called critical density

$$\rho_{\text{tot}}(\eta) = H^2(\eta)\varphi^2(\eta) \equiv \rho_{\text{cr.}}(\eta).$$

The permanent dominance of the matter with the rigid state means the existence of the integral of motion

$$H(\eta)\varphi^2(\eta) = H_0\varphi_0^2.$$

Now find a ratio of the density of the created matter $\rho_v(\eta_I) \sim T^4 \sim H_I^4 \sim M_I^4$ to the density of the primordial cosmological motion of the universe $\rho_{\text{cr.}}(\eta) = H_I^2\varphi_I^2$. This ratio has an extremely small number

$$\frac{\rho_v(\eta_I)}{\rho_{\text{cr.}}(\eta_I)} = \frac{M_I^2}{\varphi_I^2} = \frac{M_W^2}{\varphi_0^2} = y_v^2 = 10^{-34}.$$

Therefore the back reaction of created particles on the evolution of the universe is a negligible quantity. For the lifetime of the universe the primordial density of the cosmological motion $\rho_{\text{tot}}(\eta) = H_I^2\varphi_I^4/\varphi^2(\eta) \equiv H_0^2\varphi_0^4/\varphi^2(\eta)$ decreases by 10^{29} times, and in the present-day epoch the critical density $\rho_{\text{cr.}0} \equiv H_0^2\varphi_0^2 = 10^{-29}\rho_{\text{cr.}I}$ is $20 \div 30$ times greater than the density of the observed baryon matter.

D. Baryon asymmetry

The massive vector bosons during their lifetime polarize the Dirac sea vacuum of filled states of

left fermions with negative energy and create the baryon asymmetry of the universe [51].

Interaction of the primordial W and Z bosons with the left-hand fermions leads to nonconservation of fermion quantum numbers. It is known that the gauge-invariant current of each doublet is saved only at a classic level [52].

At a quantum level, we have an abnormal current $j_{\mu L}^{(i)} = \psi_L^{(i)} \gamma_\mu \psi_L^{(i)}$,

$$\partial_\mu j_{\mu L}^{(i)} = -\frac{\text{Tr} \hat{F}_{\mu\nu}^* \hat{F}_{\mu\nu}}{16\pi^2},$$

where

$$\hat{F}_{\mu\nu} = -\frac{ig\tau_a}{2}(\partial_\mu v_\nu^a - \partial_\nu v_\mu^a).$$

During their lifetime, the vector bosons polarize the Dirac sea of the left-hand fermions with negative energy. If we take the integral of the abnormal current $\partial_0 j_{0L}$ over the four-dimensional conformal space-time confined between three-dimensional hyperplanes $\eta = 0$ and $\eta = \eta_L$, we find that the number of the left-hand fermions is equal to

$$\Delta N = \frac{2\pi\alpha_{\text{QED}}}{\sin^2 \theta_W} \left[4N_W + \frac{N_Z}{\cos \theta_W} \right]$$

where $N_v(\eta_L)$ is determined by the expectation value of the Chern-Simons functional over the vacuum (87) [52]

$$N_v(\eta_L) = -\int_0^{\eta_L} d\eta \int \frac{d^3x}{16\pi^2} \text{sq} \langle 0 | \text{Tr} F_{\mu\nu}^v * F_{\mu\nu}^v | 0 \rangle_{\text{sq}},$$

where η_L is the lifetime of bosons.

If we take into account the numerical evaluation of N_W , N_Z , in the conjecture that the lifetime of bosons is $\tau_L^W = 15$, $\tau_L^Z = 30$, we estimate the magnitude of the nonconservation of the fermion number

$$\Delta F = \frac{\Delta N}{V_0} \simeq 1.2n_\gamma$$

where n_γ is the density of the number of the CMB photons. The baryon asymmetry appears as a consequence of three Sakharov conditions [52]: the CP-nonconservation, the evolution of the universe $H_0 \neq 0$ and the violation of the baryon number

$$\Delta B = X_{\text{CP}} \frac{\Delta F}{3} \simeq 0.4X_{\text{CP}}n_\gamma,$$

where X_{CP} is a factor determined by a superweak interaction of d and s -quarks ($d + s \rightarrow s + d$) with the CP-violation experimentally observed in decays of K mesons [53]. From a ratio of the number of baryons to the number of photons it is possible to make an estimation of magnitudes of a constant of a weak coupling $X_{\text{CP}} \sim 10^{-8} \div 10^{-9}$.

V. SUMMARY

A. Theory

We have shown that there are a set of theoretical and observational arguments in favour of Conformal General Relativity (CGR) described by the action

$$S_{\text{CGR}}^{(2)}[w|\chi, F] = S_{\text{PCT}}^{(2)}[w|\chi, g] + S_{\text{SM}}[w|F],$$

where $S_{\text{SM}}[w|F]$ is the Standard Model (SM) with the dilaton scalar field w instead of the Higgs mass and F is the set of fields of SM including the metric components $g_{\mu\nu}$. The dilaton w and its angle χ are described by the difference of two PCT-actions (26)

$$S_{\text{PCT}}^{(2)}[w|\chi, g] = S_{\text{PCT}}[w_-|g] - S_{\text{PCT}}[w_+|g],$$

with $w_+ = w \cosh \chi$, $w_- = w \sinh \chi$. We called the dilaton angle χ the *quintessence*. This theory is invariant with respect to a large group of the gauge, general coordinate, and conformal transformations.

B. Method

Any relativistic quantum theory requires its physical meaning only for description of concrete physical objects including our universe as a finite part of the world with a finite volume V_0 and a finite lifetime $\eta \sim 10^{17} \text{sec}$.

It is useful to recall the following words by Max Born about quantum theory ([54], p. 108): "The clue is the point ..., that quantum mechanics does not describe a situation in an objective external world, but a definite experimental arrangement for observing a section of the external world. Without this idea even the formulation of a dynamical problem in quantum theory is impossible. But if it is acceptable, the fundamental indeterminacy in the physical predictions becomes natural as no experimental arrangement can ever be absolute precise." Following to Max Born one can say that the Hamiltonian description of any quantum system is determined by "a definite experimental arrangement for observing a section of the external world". If it is acceptable, the frame-dependence of the fundamental scheme of quantization becomes natural as no experimental arrangement can ever be absolute relativistic invariant.

A frame of reference (as a experimental arrangement for observing the universe with the finite volume V_0) is defined as the choice of the evolution parameter

$$w(x^0, \mathbf{x}) = \varphi(x^0)$$

in the world space of all fields ($w|\chi, F$) with corresponding measurable quantities $F = F_c$ including an interval

$$ds_c^2 = (N_c dx^0)^2 - {}^{(3)}g_{ij}^c(dx^i + N^i dx^0)(dx^j + N^j dx^0)$$

with the constraint of the finite volume

$$V[g_c] = V_0.$$

The initial CGR action in this frame defines the invariant geometric time

$$d\eta = N_0(x^0)dx^0$$

in terms of the global lapse-function $N_0(x^0)$ determined by the Gibbs averaging operation of the local one N_c

$$N_0^{-1} = \langle N_c^{-1} \rangle \equiv \frac{\int d^3x N_c^{-1} \sqrt{{}^{(3)}g_c}}{\int d^3x \sqrt{{}^{(3)}g_c}}.$$

The Gibbs averaging of the equations of motion $\langle \delta S_{\text{CGR}} / \delta g_c \rangle = 0$ leads to the cosmological equations [14, 38]

$$\varphi'^2 = \rho; \quad 3\varphi'^2 - (\varphi^2)'' = 3p,$$

where $\rho = \langle T_0^0 \mathcal{N} \rangle$ and $3p = \langle T_k^k \mathcal{N} \rangle$ are defined by the Gibbs averaging the total energy-momentum tensor with $\mathcal{N} = N_c/N_0$.

The problem of energy and time of any relativistic system (a particle [10, 11], a string [12, 13, 14] and a universe [14, 21, 36]) is solved by the identification of the measurable Hamiltonian of evolution with respect to the evolution parameter with the canonical momentum of this parameter, in our case, with P_φ . The positive value of this momentum

$$P_\varphi = 2V_0\sqrt{\rho}$$

determines the energy of the universe created at the point $\varphi_I = \varphi(\eta)|_{\eta=0}$ with the initial data of all fields including quintessence χ .

The relativistic dynamic of the universe in the world field space with respect to the conformal time $F(\eta)$ could not be described by the naive Newton-like system. The universe has two Newton-like realities with two evolution parameters φ, η , two world spaces: the field

$$\varphi, \quad F_c = \sum [a^+ + a]$$

and geometric

$$\eta, \quad F_g = \sum [b^+ + b],$$

and two wavefunctions connected by the Levi-Civita transformation [36]. The Hubble law $\varphi(\eta)$ as the relation between two Newton-like realities (i.e, two their wavefunctions) could not be understood in the naive Wheeler-DeWitt quantum theory.

C. Results: Cold Universe Scenario

At the beginning the universe was empty. Only quintessence was in the motion. The energy density is determined by the integral of motion of the quintessence $p_\chi = 2\varphi^2\chi'$

$$\rho = \rho_{\text{rigid}} = \frac{p_\chi^2}{4\varphi^2}.$$

Due the constraint $\rho = \varphi'^2$ this rigid state of the quintessence gives an integral of motion of the dilaton

$$p_\chi = (\varphi^2)' \equiv 2\varphi_I^2 H_I \quad (H = \varphi'/\varphi),$$

as an analogy of the equation $E = mc^2$ in SR.

The next event is the intensive creation from vacuum of the massive vector bosons with masses M_I and the temperature as integral of motion of the rigid state

$$T = (M_I^2 H_I)^{1/3} = (M_0^2 H_0)^{1/3} = 2.7K$$

during the time $\eta_I = 1/2H_I \sim 10^{-12} \text{sec}$.

The next step is the formation of the baryon symmetry of the universe as a consequence of the "polarization" of the Dirac sea vacuum of left fermions by the transverse bosons, according to the selection rules of the Standard Model. This sea vacuum was introduced by Dirac to cancel the negative energies and make stable the states of quantum particles. After the decay of bosons, their temperature is inherited by the CMB radiation. All the subsequent evolution of matter with varying masses in the constant Universe is similar to the well-known scenario of the hot universe, like the Copernicus epicycles replicate the ones of Ptolemaeus.

The single rigid state of the free quintessence scalar field χ explains all eras of evolution of the universe [22].

VI. CONCLUSION

Physics is the science about the measurable part of our world. The complete set of all measurable physical quantities forms a manifold of initial data.

There are three levels of classification of initial data: phenomenological laws, equations of motion, and symmetry principles.

The determination of symmetry groups of initial data and their invariant structure relations in the form of physical laws is the guideline of modern physics. Initial data distinguish two types of symmetries: initial data are covariant with respect to a frame-transformation, and they are invariant with respect to a gauge-transformation.

The group of covariance of initial data in the Newton mechanics is the Galilei one. The group of covariance of initial data in the standard general relativity (SGR) as the sum of Einstein's theory and the SM is the Poincare group.

We listed here the set of arguments in favour of that the group of covariance of initial data in our world is the conformal group including scale transformations.

The first argument was historical: the symmetry of the world is the symmetry of the Faraday-Maxwell electrodynamics, i.e, relativistic, conformal, and gauge.

The second set of arguments were observational: latest SN data, primordial nucleosynthesis, and the CMBR temperature 2.7 are compatible with the conformal principle of the covariance of the initial data.

Why there exist the conformal and gauge symmetries? Why are certain symmetries (relativistic and conformal) broken by the initial data, while the others (gauge and general-covariant) limit these data by the constraint equations. Why there is such an exact fitting of the initial data and the dimensionless coupling constants to the anthropic principle? Perhaps, in future, somebody will find the answers to these questions in the harmony of the symmetry principles themselves, as the latter was found in epicycles by Copernicus and Kepler, and in dynamic laws by Einstein, Weyl, and other physicists of the XX th century.

One of the authors (V.N.P.) is grateful to N.A. Chernikov, Ch. Isham, T.W.B. Kibble, and W. Thirring for discussions.

-
- [1] S. Schweber, *An Introduction to Relativistic Quantum Field Theory* (Row, Peterson and Co • Evanston, Ill., Elmsford, N.Y 1961).
- [2] P.A.M. Dirac, Proc.Roy.Soc. A **114**, 243 (1927) ; Can. J. Phys.**33** , 650 (1955).
- [3] W. Heisenberg, and W. Pauli, Z.Phys. **56**, 1 (1929); Z.Phys. **59**, 166 (1930).
- [4] J. Schwinger, Phys. Rev. **127**, 324 (1962).
- [5] Nguyen Suan Han, and V.N. Pervushin, *Mod. Phys. Lett.* **A2** 367 (1987); V.N. Pervushin, *Dirac Variables in Gauge Theories, Lecture notes in DAAD Summerschool on Dense Matter in Particle - and Astrophysics, JINR, Dubna, Russia, August 20 - 31, 2001*; [hep-th/0109218]; Phys. Particles and Nuclei **34** (2003), in print.
- [6] L. Faddeev, V. Popov, Phys. Lett. B **25**, 29 (1967).
- [7] R. Feynman, Phys.Rev.**76**, 769 (1949).
- [8] R. Feynman, Acta Physica Polonica **24**, 697 (1963).
- [9] L.D. Faddeev, Teor. Mat. Fiz. **1**, 3 (1969), in Russian.
- [10] H. Poincare, C.R. Acad. Sci., Paris, **140**, 1504 (1904).
- [11] A. Einstein, Annal. d. Phys., **17**, 891 (1905).
- [12] B.M. Barbashov and N.A. Chernikov, *Classical dynamics of relativist c string* [in Russian], Preprint JINR P2-7852, Dubna (1974).
- [13] A.G. Reiman and L.D. Faddeev, Vestn. Leningr. Gos. Univ., No.1, 138 (1975).
- [14] B.M. Barbashov, V.N. Pervushin, and D.V. Proskurin, *Theoretical and Mathematical Physics* **132**, 1045 (2002) (Translated from Teoreticheskaya Matematicheskaya Fizika, Vol.132, No.2, pp.181 -197, August, 2002).
- [15] B.M. Barbashov and V.V. Nesterenko, *Introduction to Relativisti String Theory* [in Russian], Energoatomizdat, Moscow (1987); English transl., World Scientific, Singapore (1990).
- [16] V.N. Pervushin, Riv. Nuovo Cim. **8**, N **10**, 1 (1985).
- [17] A.A. Friedmann, Z. für Phys, **10**, 377 (1922); **21**, 306 (1924).
- [18] H.P. Robertson, Rev Mod Phys, **5**, 62 (1933); A.G. Walker, Monthly Notice of Royal Soc, **94**, N2, 154 (1933); **95**, N3, 263 (1935).
- [19] G. Lemaitre, Monthly Notice of Royal Soc, **91**, N5, 490 (1931); G. Ann. de la Soc. Scien. de Bruxelles, **A 47**, 49 (1927).
- [20] A. Einstein and W. de-Sitter, Proc. of Nat. Acad. of Scien., **18**, N3, 213 (1932).
- [21] M. Pawlowski, V. V. Papoyan, V. N. Pervushin and V. I. Smirichinski, Phys. Lett. B 444 (1998) 293.
- [22] D. Behnke, D.B. Blaschke, V.N. Pervushin, D.V. Proskurin, Phys. Lett. B **530**, 20 (2002); [gr-qc/0102039]
- [23] M. Pawlowski and R. Raczka, Found. of Phys. 24 (1994) 1305.
- [24] R. Kallosh, L. Kofman, A. Linde and A. Van Proeyen, Class. Quant. Grav. 17 (2000) 4269; [hep-th/0006179].
- [25] R. Penrose, *Relativity, Groups and Topology*, (Gordon and Breach, London 1964); N. Chernikov and E. Tagirov, Ann. Inst. Henri Poincarè 9, 109 (1968).
- [26] H. Weyl, Sitzungsber. d. Berl. Akad. 465 (1918).
- [27] J.V. Narlikar, Space Sci. Rev. **50**, 523 (1989).
- [28] A.G. Riess *et al.*, Astron. J. **116**, 1009 (1998); S. Perlmutter *et al.*, Astrophys. J. **517**, 565 (1999).
- [29] A.G. Riess *et al.*, Astrophys. J. **560**, 49 (2001); [astro-ph/0104455].
- [30] S. Weinberg, *First Three Minutes. A modern View of the Origin of the universe* (Basic Books, Inc., Publishers, New-York, 1977).
- [31] A.D. Linde, *Elementary Particle Physics and Inflation Cosmology*, (Nauka, Moscow, 1990), in Russian.

- [32] V.I. Ogievetsky and I.V. Polubarinov, Zh. Eksp. Teor. Fiz. **41**, 246 (1961); A.A. Slavnov and L.D. Faddeev, Teor. Mat. Fiz. **3**, 18 (1970).
- [33] H.-P. Pavel and V.N. Pervushin, Int. J. Mod. Phys. A **14**, 2285 (1999).
- [34] V.N. Pervushin and D.V. Proskurin, Gravitation and Cosmology, **8**(2002) in print.
- [35] C .W. Misner, *Phys. Rev.* **186**, 1319 (1969).
- [36] M. Pawlowski, V. N. Pervushin, Int. J. Mod. Phys. **16**, 1715 (2001), [hep-th/0006116]; V. N. Pervushin and D. V. Proskurin, Gravitation and Cosmology, **7**, 89 (2001).
- [37] V. N. Pervushin and V. I. Smirichinski, J. Phys. A: Math. Gen. **32** (1999) 6191.
- [38] V.N. Pervushin and D.V. Proskurin, gr-qc/0205084.
- [39] J.C. Maxwell, *A Treatise on Electricity and Magnetism*, Oxford, 1873.
- [40] P. A. M. Dirac, *Proc.Roy.Soc.* **A 246**, 333 (1958); P. A. M. Dirac, *Phys. Rev.* **114**, 924 (1959).
- [41] R. Arnowitt, S. Deser, and C .W. Misner, *Phys. Rev.* **116**, 1322 (1959); *Phys. Rev.* **117**, 1595 (1960); *Phys. Rev.* **122**, 997 (1961).
- [42] L. D. Faddeev and V. N. Popov, *Us.Fiz.Nauk*, **111**, 427 (1973).
- [43] A.L. Zel'manov, Doklady AN SSSR **227**, 78 (1976); Vladimirov Yu.S. *Frame of references in theory of gravitation* (Moscow, Eneroizdat, 1982), in Russian.
- [44] E. A. Tagirov, N. A. Chernikov, Preprint JINR P2-3777, Dubna, 1968.; K.A. Bronnikov, E. A. Tagirov, Preprint JINR P2-4151, Dubna, 1968.
- [45] G. L. Parker, *Phys. Rev. Lett.* **21**, 562 (1968); *Phys. Rev.* **183**, 1057 (1969); *Phys. Rev. D* **3**, 346 (1971).
- [46] A.A. Grib, S.G. Mamaev, V.M. Mostepanenko "Quantum effects in intensive external fields" (Moscow, Atomizdat, 1980) (in Russian). A.A. Grib, S.G. Mamaev, *Yadernaya Fizika*, **10**, 1276 (1969).
- [47] R. U. Sexl and H. K. Urbantke, *Phys. Rev.*, **179**, 1247 (1969).
- [48] Ya.B. Zel'dovich, Pis'ma v ZHETF **12**, 443 (1970).
- [49] Ya.B. Zel'dovich, A.A. Starobinski, ZHETF **61** (1971) 2161.
- [50] D. Blaschke *et al.*, *Cosmological creation of vector bosons and CMB*, gr-qc/0103114; V.N. Pervushin and D.V. Proskurin, *Proceeding of the V International Conference on Cosmoparticle Physics (Cosmion-2001) dedicated to 80-th Anniversary of Andrei D. Sakharov* (21-30 May 2001, Moscow-St.Peterburg, Russia) gr-qc/0106006.
- [51] D. Blaschke *et. al.*, *Yad. Fiz.* (submitted); hep-th/0206246.
- [52] A. D. Sakharov, Pis'ma v ZHETF, **5**, 24 (1967); V. A. Rubakov and M. Shaposhnikov, *Uspeki Fiz. Nauk*, **166**, 493 (1996).
- [53] L. B. Okun, *Leptons and Quarks* (Nauka, Moscow 1981).
- [54] M. Born, *Natural Philosophy of Cause and Change*, 1949, Clarendon Press, Oxford.